


Protocol Information

Dave Skinner
PMC Farm Manager
Pullman Plant Materials
Center
Room 211A Hulbert Hall
WSU
Pullman,
Washington 99163-6211

509-335-9689
509-335-2940 Fax
abbie@wsu.edu



Family Scientific Name: **Poaceae**

Family Common Name: **Grass**

Scientific Name: ***Deschampsia elongata* (Hook.)
Munro ' '**

Common Synonym: ' '

Common Name: **Slender hairgrass**

Species Code: **DEEL**

Ecotype: **near Albion, WA**

General Distribution: **Western North America from
Alaska east to Alberta and
south to New Mexico
(Hitchcock & Cronquist 1973).
Mean annual precipitation
range is from 10-24 inches
(USDA NRCS 2006). It may be
found in moist meadows and
along streams (Hitchcock &
Cronquist 1971, Mohlenbrock
undated). In eastern
Washington it may also occur
on drier soils (Piper & Beattie
1914, St. John 1963). Wetland**

indicator status is FACW- (US Fish and Wildlife Service 1988).

Known Invasiveness:

Propagation Goal: Plants

Propagation Method: Seed

Product Type: Container (plug)

Stock Type: 10 cu. in.

Target Specifications: Tight root plug in container.

Propagule Collection: Seed is collected when the inflorescence begins to dry and the seed is in the soft to hard dough stage but before it shatters from the inflorescence. Seed can be stripped from the inflorescence or the inflorescence can be clipped from the plant.

Propagule Processing: Small amounts are rubbed to free the seed, then cleaned with an air column separator. Larger amounts can be threshed with a hammermill, then cleaned with air screen equipment. Clean seed is stored in controlled conditions at 40 degrees Fahrenheit and 40% relative humidity. Seed is greyish tan in color and small.

1,860,000 seeds/lb (USDA NRCS 2006).

Pre-Planting Treatments: **Laboratory germination is best with gibberellic acid and a 5 day prechill at 5 degrees centigrade (Chirco & Turner 1986).**

Seed germinates readily without pretreatment. Trials conducted at the PMC comparing untreated seed with cold, moist stratified seed showed no benefit from stratification. Seeds given 30 days on cold, moist stratification began germinating a few days earlier than untreated seed, but took longer to reach maximum germination levels.

Growing Area Preparation/
Annual Practices for Perennial Crops:

In January seed is sown in the greenhouse in 10 cu. in. Ray Leach Super cell conetainers filled with Sunshine #1 and covered lightly. Head space of ¼ to ½ inch is maintained in conetainers to allow deep watering. A thin layer of pea gravel is applied to prevent seeds from floating. Conetainers are watered deeply.

Establishment Phase: **Medium is kept moist until germination occurs. Germination usually begins in 5 days and is complete in 7 days.**

Length of Establishment Phase: **1 week**

Active Growth Phase: **Plants are watered deeply every other day and fertilized once per week with a complete, water soluble fertilizer containing micro-nutrients.**

Length of Active Growth Phase: **3 months**

Hardening Phase: **Plants are moved to the cold frame in late March or early April, depending on weather conditions. They are watered every other day if the weather is cool, and every day during hot, dry spells.**

Length of Hardening Phase: **2-4 weeks**

Harvesting, Storage and Shipping:

Length of Storage:

Outplanting performance on typical sites: **Transplanting is done in early May by using an electric drill and portable generator to drill 1.5 inch diameter holes at the planting site. Survival in seed increase plantings without competing vegetation approaches 100%. Transplanting into sites with existing vegetation may reduce survival and vigor depending on weather conditions following planting. Plants will flower and produce seed the same season.**

Other Comments: **No insect or disease problems have been noted. Plants probably can be propagated by division. This method should only be used for plants growing in cultivation. Plants should not be dug up from stands in the wild.**

References: **Chirco, Ellen, and Terry Turnoer. 1986. Species without AOSA Testing Procedures. The Newsletter of the Association of Official Seed Analysts 60 (2):2-66. Available online at <http://www.aosaseed.com/Species%20wo%20AOSA%20list%20plus%20adds.pdf>. Updated November 11/10/03. Accessed 2/16/06.**

Hitchcock, A.S. 1971. Manual of the Grasses of the United States. 2nd edition. Revised by Agnes Chase. Dover Publications, Inc., New York, NY. 2 vol.

Hitchcock, C. Leo, and Arthur Cronquist. 1973. Flora of the Pacific Northwest. University of Washington Press. Seattle, WA. 730 pp.

Mohlenbrock, Robert H. Undated. Western Wetland Flora: A Field Office Guide to Wetland Species. USDA, NRCS Western Region. Sacramento, CA.

Piper, C.V., and R.K. Beattie. 1914. The Flora of Southeastern Washington and Adjacent Idaho. Lancaster, PA. Press of the New Era Printing Company. 296 p.

St. John, Harold. 1963. Flora of Southeastern Washington and of Adjacent Idaho. 3rd edition. Outdoor Pictures. Escondido, CA. 583 pp.

USDA, NRCS. 2006. The PLANTS Database, Version 3.5 (<http://plants.usda.gov>). Data compiled from various sources by Mark W. Skinner. National Plant Data Center, Baton Rouge, LA 70874-4490 USA.

US Fish and Wildlife Service. 1988. National list of vascular plant species that occur in wetlands. US Fish & Wildlife Service Biological Report 88 (24).

Citation:

Skinner, David M. 2006. Propagation protocol for production of container *Deschampsia elongata* (Hook.) Munro ' ' plants (10 cu. in.); Pullman Plant Materials Center, Pullman, Washington. In: Native Plant Network. URL: <http://www.nativeplantnetwork.org> (accessed 14 March 2006). Moscow (ID): University of Idaho, College of Natural Resources, Forest Research Nursery.